

Assessment of Influences of Stenoses in Right Carotid Artery on Left Carotid Artery Using Wall Stress Marker

Bit A., Ghagare D., Rizvanov A., Chattopadhyay H.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017 Arindam Bit et al. Purpose. Atherosclerosis is a diseased condition of blood vessel. It causes partial blockage in lumen of vessel and affects hemodynamic of localized flowing blood. Complex geometries like region of bifurcation also affects hemodynamic to a larger extent. Complexity further increases in presence of stenoses at region of bifurcation. Such morphological change in vessel largely affects parent as well as corresponding sister and daughter vessels. In this paper, complexity in hemodynamic of blood in pair of carotid arteries (left and right carotid arteries) is evaluated in presence of stenoses at basilar segment of right artery in three-dimensional domain using reconstructed tomographic images of patient. Methods. Transient information of blood flow is obtained using four-dimensional phase-contrast MRI technique. Haematocrit component of blood at diseased condition is considered using Power Law and Quemada model. Numerical techniques are used to solve pressure-coupled governing equations of flowing blood. Results. Dysfunctions of endothelial cells near the wall are characterised by evaluating shear stress markers. Wall shear stress and its gradient based and harmonic based descriptors are calculated over complete geometry during one cardiac cycle. Conclusion. Internal branch of left carotid artery and external branch of right carotid artery are found prone to secondary stenoses in presence of primary stenoses at basilar segment of right carotid artery.

<http://dx.doi.org/10.1155/2017/2935195>

References

- [1] D. Drikakis, C. Milionis, S. K. Pal, S. Patel, and E. Shapiro, "Assessment of the applicability of analytical models for blood flow prediction in reconstructive surgery, " *International Journal for Numerical Methods in Biomedical Engineering*, vol. 27, no. 7, pp. 993-999, 2011.
- [2] P. Neofytou and D. Drikakis, "Effects of blood models on flows through a stenosis, " *International Journal for Numerical Methods in Fluids*, vol. 43, no. 6-7, pp. 597-635, 2003.
- [3] Y. Jinyou and H. Yang, "Numerical simulations of the nonnewtonian blood flow in human thoracic aortic dissection based on CT images, " in *Proceedings of the 5th International Conference on Bioinformatics and Biomedical Engineering, (ICBBE 11)*, pp. 1-4, Wuhan, China, May 2011.
- [4] U. Morbiducci, D. Gallo, D. Massai et al., "On the importance of blood rheology for bulk flow in hemodynamic models of the carotid bifurcation, " *Journal of Biomechanics*, vol. 44, no. 13, pp. 2427-2438, 2011.
- [5] D. N. Ku, D. P. Giddens, C. K. Zarins, and S. Glagov, "Pulsatile flow and atherosclerosis in the human carotid bifurcation. Positive correlation between plaque location and low and oscillating shear stress, " *Arteriosclerosis*, vol. 5, no. 3, pp. 293-302, 1985.
- [6] A. C. Benim, A. Nahavandi, A. Assmann, D. Schubert, P. Feindt, and S. H. Suh, "Simulation of blood flow in human aorta with emphasis on outlet boundary conditions, " *Applied Mathematical Modelling. Simulation and Computation for Engineering and Environmental Systems*, vol. 35, no. 7, pp. 3175-3188, 2011.

- [7] K. Perktold, M. Resch, and H. Florian, "Pulsatile non-Newtonian flow characteristics in a three-dimensional human carotid bifurcation model, " *Journal of Biomechanical Engineering*, vol. 113, no. 4, pp. 464-475, 1991.
- [8] F. P. P. Tan, G. Soloperto, S. Bashford et al., "Analysis of flow disturbance in a stenosed carotid artery bifurcation using two-equation transitional and turbulence models, " *Journal of Biomechanical Engineering*, vol. 130, no. 6, Article ID 061008, 12 pages, 2008.
- [9] J. B. Thomas, L. Antiga, S. L. Che et al., "Variation in the carotid bifurcation geometry of young versus older adults: implications for geometric risk of atherosclerosis, " *Stroke*, vol. 36, no. 11, pp. 2450-2456, 2005.
- [10] L. Grinberg, T. Anor, J. R. Madsen, A. Yakhot, and G. E. Karniadakis, "Large-scale simulation of the human arterial tree, " *Clinical and Experimental Pharmacology and Physiology*, vol. 36, no. 2, pp. 194-205, 2009.
- [11] D. Gallo, D. A. Steinman, P. B. Bijari, and U. Morbiducci, "Helical flow in carotid bifurcation as surrogate marker of exposure to disturbed shear, " *Journal of Biomechanics*, vol. 45, no. 14, pp. 2398-2404, 2012.
- [12] S. R. Shah, "Capillary-Tissue diffusion phenomena for blood flow through a stenosed artery using Herschel-Bulkley fluid, " *International Journal of Biochemistry and Biophysics*, vol. 1, pp. 1-8, 2011.
- [13] F. J. Walburn and D. J. Schneck, "A constitutive equation for whole human blood, " *Biorheology*, vol. 13, no. 3, pp. 201-210, 1976.
- [14] S. Chien, S. Usami, R. J. Dellenback, and M. I. Gregersen, "Blood viscosity: influence of erythrocyte deformation, " *Science*, vol. 157, no. 3790, pp. 827-829, 1967.
- [15] D. Quemada, "Rheology of concentrated disperse systems III. General features of the proposed non-newtonian model. Comparison with experimental data, " *Rheologica Acta*, vol. 17, no. 6, pp. 643-653, 1978.
- [16] S. V. Patankar, *Numerical Heat Transfer and Fluid Flow*, McGraw-Hill, New York, NY, USA, 1980.
- [17] C. M. Rhie and W. L. Chow, "Numerical study of the turbulent flow past an airfoil with trailing edge separation, " *AIAA Journal*, vol. 21, no. 11, pp. 1525-1532, 1983.
- [18] T. Nandi and H. Chattopadhyay, "Simultaneously developing flow in microchannels under pulsating inlet flow condition, " *International Journal of Transport Phenomena*, vol. 13, pp. 110-120, 2012.
- [19] C. Huang, Z. Chai, and B. Shi, "Non-newtonian effect on hemodynamic characteristics of blood flow in stented cerebral aneurysm, " *Communications in Computational Physics*, vol. 13, no. 3, pp. 916-928, 2013.
- [20] V. Deplano and M. Siouffi, "Experimental and numerical study of pulsatile flows through stenosis: wall shear stress analysis, " *Journal of Biomechanics*, vol. 32, no. 10, pp. 1081-1090, 1999.
- [21] H. A. Himburg, D. M. Grzybowski, A. L. Hazel, J. A. LaMack, X.-M. Li, and M. H. Friedman, "Spatial comparison between wall shear stress measures and porcine arterial endothelial permeability, " *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 286, no. 5, pp. H1916-H1922, 2004.
- [22] A. Bit and H. Chattopadhyay, "Numerical investigations of pulsatile flow in stenosed artery, " *Acta of Bioengineering and Biomechanics*, vol. 16, no. 4, pp. 33-44, 2014.